

Section 1

Tissue and Structure mcqs

- 1) Regarding tissues:
 - a) cartilage is very vascular
 - b) ligaments are generally not elastic
 - c) deep fascia is insensate
 - d) cardiac muscle is not striated
 - e) periosteum is not sensitive

- 2) An example of a unipennate muscle is:
 - a) sartorius
 - b) flexor pollicis longus
 - c) rectus femoris
 - d) deltoid
 - e) tibialis anterior

- 3) An example of a secondary cartilaginous joint is:
 - a) distal tibiofibular joint
 - b) costochondral joint
 - c) sagittal suture
 - d) manubrio-sternal joint
 - e) knee joint

- 4) Which vessel is not an end artery?
 - a) renal artery
 - b) retinal artery
 - c) pulmonary artery
 - d) intercostal artery
 - e) splenic artery

- 5) The appendages of the skin:
 - a) are derived from mesoderm
 - b) are derived from epidermal tissues
 - c) lie in the dermal layer
 - d) are derived from endoderm

- 6) Melanocyte(s):
 - a) number determine the colour of the skin
 - b) are responsible for high colour, greying is the result of decreasing numbers
 - c) produce varying melanins
 - d) are found mainly in the dermal layer

- 7) In regards to connective tissue:
 - a) aponeuroses are a form of deep fascia
 - b) retinaculae are a form of ligament
 - c) deep fascia is insensate
 - d) some ligaments are designed to allow a degree of elasticity

- 8) With regards to cartilage:
- a) in the elderly, the epiglottis can occasionally be seen on x-ray as its elastic cartilage calcifies
 - b) bones that develop by intramembranous ossification have fibrocartilage at their articular surfaces
 - c) cartilage is nourished exclusively by ground substance as it has no blood supply
 - d) the most common form of cartilage is fibrocartilage
- 9) Which is correct?
- a) oblique alignment of striated muscle fibres allows for greater force of contraction
 - b) smooth muscle is multi-nucleated
 - c) the epimysium surrounds groups of striated muscle fibres to form muscle bundles or fasciculi
 - d) rectus femoris fibres are arranged in a multi-pennate form to allow increased force of contraction
 - e) alpha anterior horn cells of the spinal cord supply the intrafusal muscle fibres
- 10) Which is correct?
- a) B-cells are responsible for cell-mediated immunity
 - b) plasma cells are differentiated B-cells
 - c) all lymphoid tissue is encapsulated
 - d) in lymph nodes, follicles of lymphocytes are concentrated in the medullary region
- 11) Which of the following pairs do not match with regard to mode of ossification?
- a) ribs and vertebra
 - b) clavicle and humerus
 - c) femur and 1st metatarsal
 - d) patella and pisiform
 - e) mandible and clavicle
- 12) Regarding cartilage, which is INCORRECT?
- a) it is essentially avascular
 - b) hyaline cartilage contains hyaluronic acid
 - c) rib cartilage is elastic type
 - d) TMJ is fibrous
 - e) all contain mucopolysaccharides
- 13) Regarding muscle strength, which is not a factor?
- a) resting muscle length
 - b) cross sectional area
 - c) lever arm length
 - d) the extent to which the muscle is contracted
 - e) configuration of fibres
- 14) Which is true?
- a) epidermis is derived from mesoderm
 - b) the sweat glands, sebaceous glands are mesodermal in origin
 - c) the epidermis contains fibrous tissue and blood vessels
 - d) the colour of the skin is determined by the number of melanocytes
 - e) the dermis has a mesodermal origin

- 15) Regarding sweat glands, which is true?
- the main nerve supply is via cholinergic sympathetic fibres
 - the main nerve supply is via adrenergic sympathetic nerves
 - they are derived from mesoderm
 - the normal tubular sweat glands undergo cyclical changes with the menstrual cycle
 - the highest number of sweat glands are found on the feet
- 16) Regarding cartilage, which is true?
- they have a rich blood supply
 - the intervertebral discs contain hyaline cartilage
 - hyaline and fibrous cartilage never calcify
 - elastic cartilage never calcifies
 - elastic cartilage is the commonest type
- 17) Regarding muscles, which is true?
- cardiac muscle is more powerful than skeletal or smooth
 - skeletal muscle fibres are spindle shaped
 - smooth muscle does not contain actin or myosin
 - skeletal muscle contraction causes peristalsis in the GI tract and ureter
 - cardiac muscle fibres are short, branched, mononuclear and striated
- 18) Regarding periosteum, which is TRUE?
- it is avascular
 - it is osteogenic
 - it covers the articular surface of long bones
 - it covers the inner surface of the medullary cavity
 - it receives its blood supply from overlying muscle
- 19) An example of secondary cartilaginous joint is:
- hip joint
 - manubrio-sternal joint
 - costochondral joint
 - epiphysis
 - wrist joint
- 20) An example of a fibrous joint is:
- the structures of the skull
 - vertebral column
 - pubic symphysis
 - hip joint
 - manubrio-sternal joint
- 21) In the skin:
- apocrine sweat glands are confined to the axillae, areolar, periumbilical and genital regions
 - dark skinned races possess greater numbers of melanocytes
 - sebaceous glands are abundant on the palms and soles
 - melanocytes predominate in the dermis
 - sweat glands are under the influence of sex hormones

- 22) Regarding bone:
- a) cancellous and compact bone show similar microscopic structure
 - b) the clavicle ossifies in membrane
 - c) the bones of the skull and face ossify in cartilage
 - d) the epiphyseal plates of growing bones are composed of fibrocartilage
 - e) sesamoid bones are only associated with flexor tendons
- 23) What are vincula?
- a) the papillary ridges that form finger prints
 - b) superficial fibres of the palmar aponeurosis that insert into the skin
 - c) remnant fibres of palmar interossei that insert into the proximal area of the thumb
 - d) a communication between common flexor sheath and the sheath of pollicis longus present in 50% of people
 - e) vascular synovial folds of flexor tendons

Section 1

Tissue and Structure – Answers

- | | |
|----|-------|
| 1 | B |
| 2 | B |
| 3 | D |
| 4 | D |
| 5 | B |
| 6 | C |
| 7 | D |
| 8 | B & C |
| 9 | A |
| 10 | B |
| 11 | B |
| 12 | C |
| 13 | A |
| 14 | E |
| 15 | A |
| 16 | D |
| 17 | E |
| 18 | B |
| 19 | B |
| 20 | A |
| 21 | A |
| 22 | B |
| 23 | E |

Section 2

Nerve Anatomy

- 1) A group C nerve fibre:
 - a) is up to 20um in diameter
 - b) is myelinated
 - c) has motor function
 - d) is fusimotor to muscle spindles
 - e) is postganglionic autonomic

- 2) Regarding autonomic nervous system, visceral pain is not referred to L3-S1 because:
 - a) there are no visceral afferents related to these lumbar splanchnic nerves
 - b) there are no white rami communications to these spinal nerves
 - c) there are no sympathetic ganglia associated with these spinal segments
 - d) there is no sympathetic efferent supply to these spinal nerves
 - e) there are no grey rami communicants associated with the autonomic ganglia of L3-S1 spinal nerves

- 3) Which of the following peripheral fibres have pain and temperature fibres?
 - a) alpha neuron
 - b) beta neuron
 - c) gamma neuron
 - d) delta neuron
 - e) group B fibres

- 4) Regarding the sympathetic nervous system:
 - a) every spinal nerve receives a white ramus communicans
 - b) efferent post ganglionic fibres are myelinated
 - c) afferent sympathetic fibres synapse in the sympathetic trunk
 - d) preganglionic cell bodies lie within the lateral horn of the T1-L2 spinal segments
 - e) each cervical ganglion gives off one ramus communicans

- 5) Which of the following nerve roots is correctly associated with the corresponding muscle action?
 - a) L5 plantar flexion
 - b) L2 knee extension
 - c) C5 abduction of the shoulder
 - d) C8 abduction of the thumb
 - e) S1 extension of the great toe

Section 2

Nerve Anatomy – Answers

- | | |
|---|---|
| 1 | E |
| 2 | B |
| 3 | D |
| 4 | D |
| 5 | C |

Section 3

Nervous System

- 1) Which has the widest fibre diameter on average?
 - a) muscle spindle nerve fibres
 - b) group b
 - c) δ of group A
 - d) motor
 - e) postganglionic autonomic

- 2) Which is correct?
 - a) the lowest spinal nerve to give cutaneous supply to the anterior abdominal wall is L2
 - b) no post. ramus ever supplies skin or muscle of a limb
 - c) the posterior rami supply the prevertebral flexor muscles
 - d) C1-2 and T1 have no cutaneous branches
 - e) L1-S1 contain autonomic fibres

- 3) Which is the correct tendon reflex?
 - a) biceps C5
 - b) triceps C8
 - c) knee L3
 - d) ankle S3

Section 3

Nervous System – Answers

- | | |
|---|---|
| 1 | D |
| 2 | B |
| 3 | C |

Section 4

Embryology

- 1) At birth:
 - a) the transverse diameter of the thorax is thrice the AP diameter
 - b) the liver is relatively twice as big as that of the adult
 - c) the suprarenal gland is twice as big as the kidney
 - d) the spinal cord extends to L5
 - e) the rib cage is more vertically inclined than in the adult

- 2) In the foetal skull:
 - a) the vertical height of the orbit is equal to the combined heights of the maxilla and the mandible
 - b) the mandible is ossified at birth
 - c) the anterior fontanelle is closed at the end of one year of life
 - d) the volume of the vault is equal to that of the face
 - e) the bones of the face and vault ossify at 6 years

Section 4

Embryology – Answers

- | | |
|---|---|
| 1 | B |
| 2 | A |

Tissues and Structures

1. Regarding the ossification centres of the bones of the foot, which is INCORRECT?

- a. There are three at birth
- b. Fifth metatarsus has three ossification centres
- c. Metatarsals have two centres
- d. ?
- e. ?

2. Regarding the anatomy of the newborn

- a. The liver is relatively small
- b. The bladder lies within the pelvis
- c. The facial skeleton is relatively small
- d. The spinal cord extends to the level of the L4 vertebra
- e. The maxillary sinus is well developed at birth

3. An example of a secondary cartilaginous joint is

- a. Proximal tibial epiphysis
- b. Costochondral joint
- c. Intervertebral disc
- d. Lambdoid suture
- e. Temporomandibular joint

4. An example of a synovial joint is

- a. Intervertebral disc
- b. Sternomanubrial joint
- c. Sacroiliac joint
- d. Epiphyses
- e. Distal tibiofibular joint

Answers

- 1. b
- 2. c
- 3. c
- 4. c

1 The following are features of typical synovial joints EXCEPT

- A Bone ends are covered in hyaline or fibrous cartilage
- B A capsule encloses the joint cavity
- C The capsule is reinforced by ligaments
- D The capsule is lined by synovial membrane
- E The joint is capable of movement

2 Regarding cartilage

- A Hyaline cartilage has an effective blood supply
- B Hyaline cartilage does not have a tendency to calcify
- C The epiglottis is composed of elastic cartilage
- D The larynx is composed of fibrocartilage
- E The menisci of the knee are hyaline cartilage

3 Regarding skin

- A Skin appendages are mesodermal in origin
- B Skin colour is determined by melanocyte number
- C The greatest concentration of sweat glands is on the lip margins
- D Apocrine glands are confined to the axilla
- E There are no sebaceous glands on the palms of the hands

4 Which of the following is a primary cartilaginous joint

- A Manubriosternal joint
- B 1st sternocostal joint
- C Acromioclavicular joint
- D Inferior tibiofibular joint
- E 2nd costochondral joint

5 Which of the following is a fibrous joint

- A Pubic symphysis
- B Xiphisternal joint
- C 1st sternocostal joint
- D Superior tibiofibular joint
- E Inferior tibiofibular joint

Answers

- 1 A
- 2 C
- 3 E
- 4 B
- 5 E

24) Regarding tissues:

- a) cartilage is very vascular
- b) ligaments are generally not elastic
- c) deep fascia is insensate
- d) cardiac muscle is not striated
- e) periosteum is not sensitive

25) An example of a unipennate muscle is:

- a) sartorius
- b) flexor pollicis longus
- c) rectus femoris
- d) deltoid
- e) tibialis anterior

26) An example of a secondary cartilaginous joint is:

- a) distal tibiofibular joint
- b) costochondral joint
- c) sagittal suture
- d) manubrio-sternal joint
- e) knee joint

27) Which vessel is not an end artery?

- a) renal artery
- b) retinal artery
- c) pulmonary artery
- d) intercostal artery
- e) splenic artery

28) The appendages of the skin:

- a) are derived from mesoderm
- b) are derived from epidermal tissues
- c) lie in the dermal layer
- d) are derived from endoderm

29) Melanocyte(s):

- a) number determine the colour of the skin
- b) are responsible for high colour, greying is the result of decreasing numbers
- c) produce varying melanins
- d) are found mainly in the dermal layer

30) In regards to connective tissue:

- a) aponeuroses are a form of deep fascia
- b) retinaculae are a form of ligament
- c) deep fascia is insensate
- d) some ligaments are designed to allow a degree of elasticity

31) With regards to cartilage:

- a) in the elderly, the epiglottis can occasionally be seen on x-ray as its elastic cartilage calcifies
- b) bones that develop by intramembranous ossification have fibrocartilage at their articular surfaces
- c) cartilage is nourished exclusively by ground substance as it has no blood supply
- d) the most common form of cartilage is fibrocartilage

32) Which is correct?

- a) oblique alignment of striated muscle fibres allows for greater force of contraction
- b) smooth muscle is multi-nucleated
- c) the epimysium surrounds groups of striated muscle fibres to form muscle bundles or fasciculi
- d) rectus femoris fibres are arranged in a multi-pennate form to allow increased force of contraction
- e) alpha anterior horn cells of the spinal cord supply the intrafusal muscle fibres

33) Which is correct?

- a) B-cells are responsible for cell-mediated immunity
- b) plasma cells are differentiated B-cells
- c) all lymphoid tissue is encapsulated
- d) in lymph nodes, follicles of lymphocytes are concentrated in the medullary region

34) Which of the following pairs do not match with regard to mode of ossification?

- a) ribs and vertebra
- b) clavicle and humerus
- c) femur and 1st metatarsal

- d) patella and pisiform
 - e) mandible and clavicle
- 35) Regarding cartilage, which is INCORRECT?
- a) it is essentially avascular
 - b) hyaline cartilage contains hyaluronic acid
 - c) rib cartilage is elastic type
 - d) TMJ is fibrous
 - e) all contain mucopolysaccharides
- 36) Regarding muscle strength, which is not a factor?
- a) resting muscle length
 - b) cross sectional area
 - c) lever arm length
 - d) the extent to which the muscle is contracted
 - e) configuration of fibres
- 37) Which is true?
- a) epidermis is derived from mesoderm
 - b) the sweat glands, sebaceous glands are mesodermal in origin
 - c) the epidermis contains fibrous tissue and blood vessels
 - d) the colour of the skin is determined by the number of melanocytes
 - e) the dermis has a mesodermal origin
- 38) Regarding sweat glands, which is true?
- a) the main nerve supply is via cholinergic sympathetic fibres
 - b) the main nerve supply is via adrenergic sympathetic nerves
 - c) they are derived from mesoderm
 - d) the normal tubular sweat glands undergo cyclical changes with the menstrual cycle
 - e) the highest number of sweat glands are found on the feet
- 39) Regarding cartilage, which is true?
- a) they have a rich blood supply
 - b) the intervertebral discs contain hyaline cartilage
 - c) hyaline and fibrous cartilage never calcify
 - d) elastic cartilage never calcifies
 - e) elastic cartilage is the commonest type
- 40) Regarding muscles, which is true?
- a) cardiac muscle is more powerful than skeletal or smooth
 - b) skeletal muscle fibres are spindle shaped
 - c) smooth muscle does not contain actin or myosin
 - d) skeletal muscle contraction causes peristalsis in the GI tract and ureter
 - e) cardiac muscle fibres are short, branched, mononuclear and striated
- 41) Regarding periosteum, which is TRUE?
- a) it is avascular
 - b) it is osteogenic
 - c) it covers the articular surface of long bones

- d) it covers the inner surface of the medullary cavity
 - e) it receives its blood supply from overlying muscle
- 42) An example of secondary cartilaginous joint is:
- a) hip joint
 - b) manubrio-sternal joint
 - c) costochondral joint
 - d) epiphysis
 - e) wrist joint
- 43) An example of a fibrous joint is:
- a) the structures of the skull
 - b) vertebral column
 - c) pubic symphysis
 - d) hip joint
 - e) manubrio-sternal joint
- 44) In the skin:
- a) apocrine sweat glands are confined to the axillae, areolar, periumbilical and genital regions
 - b) dark skinned races possess greater numbers of melanocytes
 - c) sebaceous glands are abundant on the palms and soles
 - d) melanocytes predominate in the dermis
 - e) sweat glands are under the influence of sex hormones
- 45) Regarding bone:
- a) cancellous and compact bone show similar microscopic structure
 - b) the clavicle ossifies in membrane
 - c) the bones of the skull and face ossify in cartilage
 - d) the epiphyseal plates of growing bones are composed of fibrocartilage
 - e) sesamoid bones are only associated with flexor tendons
- 46) What are vincula?
- a) the papillary ridges that form finger prints
 - b) superficial fibres of the palmar aponeurosis that insert into the skin
 - c) remnant fibres of palmar interossei that insert into the proximal area of the thumb
 - d) a communication between common flexor sheath and the sheath of pollicis longus present in 50% of people
 - e) vascular synovial folds of flexor tendons

Answers

- 24 B
- 25 B
- 26 D
- 27 D
- 28 B
- 29 C
- 30 D
- 31 B & C
- 32 A

- 33 B
- 34 B
- 35 C
- 36 A
- 37 E
- 38 A
- 39 D
- 40 E
- 41 B
- 42 B
- 43 A
- 44 A
- 45 B
- 46 E

- 6) A group C nerve fibre:
- a) is up to 20um in diameter
 - b) is myelinated
 - c) has motor function
 - d) is fusimotor to muscle spindles
 - e) is postganglionic autonomic
- 7) Regarding autonomic nervous system, visceral pain is not referred to L3-S1 because:
- a) there are no visceral afferents related to these lumbar splanchnic nerves
 - b) there are no white rami communications to these spinal nerves
 - c) there are no sympathetic ganglia associated with these spinal segments
 - d) there is no sympathetic efferent supply to these spinal nerves
 - e) there are no grey rami communicans associated with the autonomic ganglia of L3-S1 spinal nerves
- 8) Which of the following peripheral fibres have pain and temperature fibres?
- a) alpha neuron
 - b) beta neuron
 - c) gamma neuron
 - d) delta neuron
 - e) group B fibres
- 9) Regarding the sympathetic nervous system:
- a) every spinal nerve receives a white ramus communicans
 - b) efferent post ganglionic fibres are myelinated
 - c) afferent sympathetic fibres synapse in the sympathetic trunk
 - d) preganglionic cell bodies lie within the lateral horn of the T1-L2 spinal segments
 - e) each cervical ganglion gives off one ramus communicans
- 10) Which of the following nerve roots is correctly associated with the corresponding muscle action?
- a) L5 plantar flexion
 - b) L2 knee extension
 - c) C5 abduction of the shoulder
 - d) C8 abduction of the thumb
 - e) S1 extension of the great toe

Answers

- 6 E
- 7 B
- 8 D
- 9 D
- 10 C

4) Which has the widest fibre diameter on average?

- a) muscle spindle nerve fibres
- b) group b
- c) δ of group A
- d) motor
- e) postganglionic autonomic

5) Which is correct?

- a) the lowest spinal nerve to give cutaneous supply to the anterior abdominal wall is L2
- b) no post. ramus ever supplies skin or muscle of a limb
- c) the posterior rami supply the prevertebral flexor muscles
- d) C1-2 and T1 have no cutaneous branches
- e) L1-S1 contain autonomic fibres

6) Which is the correct tendon reflex?

- a) biceps C5
- b) triceps C8
- c) knee L3
- d) ankle S3

Answers

- 4 D
- 5 B
- 6 C

3) At birth:

- a) the transverse diameter of the thorax is thrice the AP diameter
- b) the liver is relatively twice as big as that of the adult
- c) the suprarenal gland is twice as big as the kidney
- d) the spinal cord extends to L5
- e) the rib cage is more vertically inclined than in the adult

4) In the foetal skull:

- a) the vertical height of the orbit is equal to the combined heights of the maxilla and the mandible
- b) the mandible is ossified at birth
- c) the anterior fontanelle is closed at the end of one year of life
- d) the volume of the vault is equal to that of the face
- e) the bones of the face and vault ossify at 6 years

Answers

- 1 B
- 2 A